







- □□□□□□□□ □□□□ □□□□

□□□□

- <http://www.eu-patient.eu/globalassets/policy/adherence-compliance-concordance/adherence-joint-briefing-paper.pdf>
- [http://www.jhasim.com/files/articlefiles/pdf/ASM\\_6\\_7C\\_652-658\\_R1.pdf](http://www.jhasim.com/files/articlefiles/pdf/ASM_6_7C_652-658_R1.pdf)

□□□□

1. McHorney CA. The Adherence Estimator:A brief proximal screener for propensity to adhere to medications for chronic disease.CMR0, 25(1), 2009, 215-38.
2. g. Gardner ME, Rulien N, McGhan WF, Mead RA.A trial of patients' perceived importance of medication information provided by physicians in a health maintenance organisation.Drug Intell Clin Pharm.1988;22:596-598; Makoul G, Arntson P, Schofield T. Health promotion in primary care:physician-patient communication and decision making about prescription medications.Soc Sci Med.1995;41:1241-1254; Tarn DM, Heritage J, Paterniti DA, Hays RD, Kravitz RL, Wenger NS.Physician communication when prescribing new medications.Arch Intern Med.2006;166:1855-1862.
3. Bailey BJ, Carney SL, Gillies AH, McColm LM, Smith AJ, Taylor M. Hypertension treatment compliance:what do patients want to know about their medications?Prog Cardiovasc Nurs.1997;12:23-28; Ziegler DK, Mosier MC, Buenaver M, Okuyemi K. How much information about adverse effects of medication do patients want from physicians?Arch Intern Med.2001;161:706-713.
4. Lapane KL, Dube CE, Schneider KL, Quilliam BJ.Misperceptions of patients vs providers regarding medication-related communication issues.Am J Manag Care.2007;13:613-618.

A2-5.22-V1.0